

ALL PATIENTS ANTICIPATING SURGERY MUST STOP THE USE OF ALL SOURCES OF ASPIRIN. ASPIRIN IS A VERY STRONG ANTI-COAGULANT WHICH CAUSES PROFOUND BLEEDING PROBLEMS IN NORMAL INDIVIDUALS. THEREFORE, YOU MUST STOP TAKING ASPIRIN AND ALL ASPIRIN CONTAINING PRODUCTS

***FOR 4 WEEKS BEFORE SURGERY AND 4 WEEKS AFTER SURGERY.**

THE FOLLOWING ARE ONLY A FEW ASPIRIN CONTAINING COMPOUNDS TO BE AVOIDED:

ADVIL	COPE	NUPRIN
ALEVE	CORICIDIN	PERCODAN
ALKA SELTZER	DARVON COMPOUND	PERCOGESIC
ANACIN	DRISTAN	PABIRIN TABS
ANAPROX	DURAGESIC	PANALGESIC
A.P.C.	ECOTRIN	PERSISTIN
ASCODEEN-30	EMPRAZIL	ROBAXISAL
ASCRIPTIN	EMPIRIN	SINE-AID
ASPIRIN	EQUAGESIC	SINE-OFF
ASPIRIN	SUPPOSITORIES EXCEDRIN	SK-65 COMPOUND
BAYER ASPIRIN	FELDENE	STENDIN
BC POWDERS	FIORINAL	STERO-DARVON W/ASP.
BUFF-A-COMP	INDOCIN	SUPAC
BUFFADYNE	MEASURIN	SYNALOGOS D.C.
BUFFERIN	MIDOL	SYNALOGOS CAPSULES
BUTALBITAL	MONACET W/CODEINE	TOLECTIN
CAMA-INLAY TABS	MOTRIN	TRIAMINICIN
CHERACOL CAPSULES	NAPROSYN	VANQUICH
CLINORIL	NORGESIC	ZOMAX
CONGESPIRIN	EXCEDRIN	PEPTO BISMOL
	EXCEDRIN PM	

IF YOU MUST TAKE SOMETHING FOR HEADACHE, MENSTRUAL CRAMPS, OR OTHER ACHES AND PAINS, YOU MAY ONLY TAKE TYLENOL (AS DIRECTED) FOR THE 4 WEEKS PRIOR TO AND AFTER YOUR SURGERY.

TOPICAL ASPIRIN CONTAINING PRODUCTS TO BE AVOIDED:

ABSORBANT JR.	BRASKA	NEURABALM
ABSORBINE ARTHRITIC	COUNTERPAIN RUB	OIL-O-SOL
ABSORBINE JR.	DENCORUB	OMEGA OIL
ACT ON RUB	DOAN'S RUB	PANALGESIC
ANALBALM	EMUL-O-BALM	RID-A-PAIN
ANALGESIC BALM	END-AKE	RUMARUB
ANTIPHLOGISTINE	EXOCAINE PLUS	SLOAN'S
ANTHRALGEN	EXOCAINE TUBE	SOLTICE HI-THER
ASPERCREME	HEET	SOLTICE QUICK RUB
BANALG	ICY HOT	SPD
BAUMODYHNE	INFRA-RUB	STIMURUB
BEN GAY (ALL)	LINI-BALM	SURIN
ZEMO OINTMENT	MENTHOLATUM	YAEGER'S LINIMENT

***PLEASE DISCONTINUE ALL HOMEOPATHIC MEDICATIONS I.E., VITAMIN E., FISH OIL, OMEGA 3, GARLIC, ST,JOHNS WORT, ECHINACEA, EPHEDRINE, etc.,**